arXiv:math/9808022v1 [math.QA] 5 Aug 1998

A functional-analytic theory of vertex (operator) algebras, I

Yi-Zhi Huang

Abstract

This paper is the first in a series of papers developing a functional-analytic theory of vertex (operator) algebras and their representations. For an arbitrary $\mathbb{Z}$-graded finitely-generated vertex algebra $(V, Y, \mathbf{1})$ satisfying the standard grading-restriction axioms, a locally convex topological completion H of V is constructed. By the geometric interpretation of vertex (operator) algebras, there is a canonical linear map from $V \otimes V$ to $\overline{V}$ (the algebraic completion of V) realizing linearly the conformal equivalence class of a genus-zero Riemann surface with analytically parametrized boundary obtained by deleting two ordered disjoint disks from the unit disk and by giving the obvious parametrizations to the boundary components. We extend such a linear map to a linear map from $H\tilde{\otimes}H$ ($\tilde{\otimes}$ being the completed tensor product) to H, and prove the continuity of the extension. For any finitely-generated $\mathbb{C}$-graded V-module (W, Y_W) satisfying the standard grading-restriction axioms, the same method also gives a topological completion H^W of W and gives the continuous extensions from $H\tilde{\otimes}H^W$ to H^W of the linear maps from $V \otimes W$ to $\overline{W}$ realizing linearly the above conformal equivalence classes of the genus-zero Riemann surfaces with analytically parametrized boundaries.

0 Introduction

We begin a systematic study of the functional-analytic structure of vertex (operator) algebras and their representations in this paper.

Vertex (operator) algebras were introduced rigorously in mathematics by Borcherds and by Frenkel, Lepowsky and Meurman (see [B], [FLM] and

[FHL]). Incorporating modules and intertwining operators, the author introduced intertwining operator algebras in [H5]. The original definition of vertex (operator) algebra is purely algebraic, but a geometric interpretation of vertex operators motivated by the path integral picture in string theory was soon observed mathematically by Frenkel [F]. In [H1]–[H7], the author developed a geometric theory of vertex operator algebras and intertwining operator algebras; the hard parts deal with the Virasoro algebra, the central charge, the interaction between the Virasoro algebra and vertex operators (or intertwining operators), and the monodromies. But in this geometric theory, the linear maps associated to elements of certain moduli spaces of punctured spheres with local coordinates (or to elements of the vector bundles forming partial operads called "genus-zero modular functors" over these moduli spaces) are from the tensor powers of a vertex operator algebra (or an intertwining operator algebra) to the algebraic completion of the algebra.

To be more specific, let $(V, Y, \mathbf{1})$ be a $\mathbb{Z}$-graded vertex algebra. For any nonzero complex number z, let $P(z)$ be the the conformal equivalence class of the sphere $\mathbb{C}\cup\{\infty\}$ with the negatively oriented puncture ∞, the ordered positively oriented z and 0, and with the standard local coordinates. Then associated to $P(z)$ is the linear map $Y(\cdot, z)\cdot : V\otimes V\to \overline{V}$ given by the vertex operator map. Note that the image is in the algebraic completion $\overline{V}$ of V, not V itself.

One of the main goals of the theory of vertex operator algebras is to construct conformal field theories in the sense of Segal [S1] [S2] from vertex operator algebras and their representations, and to study geometry using the theory of vertex operator algebras. It is therefore necessary to construct locally convex topological completions of the underlying vector spaces of vertex (operator) algebras and their modules, such that associated to any Riemann surface with boundary, one can construct continuous linear maps between the tensor powers of these completions. In particular, it is necessary to construct a locally convex completion H of the underlying vector space V of a vertex algebra $(V, Y, \mathbf{1})$ such that associated to the conformal equivalence class of a disk with two smaller ordered disks deleted and with the obvious boundary parametrization, there is a canonical continuous map from $H\tilde{\otimes}H$ ($\tilde{\otimes}$ being the completed tensor product) to H. Though in some algebraic applications of conformal field theories, only the algebraic structure of conformal field theories is needed, in many geometric applications, it is necessary to have a complete locally convex topological space and continuous linear maps, associated to Riemann surfaces with boundaries, between tensor powers of the space.

In the present paper (Part I), we construct a locally convex completion H of an arbitrary finitely-generated $\mathbb{Z}$-graded vertex algebra $(V, Y, \mathbf{1})$ satisfying the standard grading-restriction axioms. The completion H is the strict inductive limit of a sequence of complete locally convex spaces constructed from the correlation functions of the generators. The strong topology, rather than the weak-$*$ topology, on the topological dual spaces of certain function spaces is needed in the construction. For any positive number r_1, r_2 and nonzero complex number z satisfying $r_2 + 2r_1 < 1$ and $r_2 < |z| < 1$, there is a unique genus-zero Riemann surface with analytically parametrized boundary given by deleting two ordered disjoint disks, the first centered at z with radius r_1 and the second centered at 0 with radius r_2, from the unit disk and by giving the obvious parametrizations to the boundary components. Associated to this genus-zero Riemann surface with analytically parametrized boundary is a linear map $Y(r_1^{L(0)}\cdot, z)r_2^{L(0)} : V \otimes V \to \overline{V}$. We extend this linear map to a linear map from $H\tilde{\otimes}H$ ($\tilde{\otimes}$ being the completed tensor product) to H, and prove the continuity of the extension. It is clear that H is linearly isomorphic to a subspace of $\overline{V}$ containing both V and the image of $Y(r_1^{L(0)}\cdot, z)r_2^{L(0)}$.

For any finitely-generated $\mathbb{C}$-graded module (W, Y_W) satisfying the standard grading-restriction axioms for such a finitely-generated vertex algebra, we also construct a locally convex completion H^W of W, and construct continuous extensions from $H\tilde{\otimes}H^W$ to H^W of the linear map $Y_W(r_1^{L(0)}\cdot, z)r_2^{L(0)} : V \otimes W \to \overline{W}$.

Our method depends only on the axiomatic properties or "world-sheet geometry" (mainly the duality properties) of the vertex algebra. Since our construction does not use any additional structure, we expect that the locally convex completions constructed in the present paper will be useful in solving purely algebraic problems in the representation theory of finitely-generated vertex (operator) algebras.

This paper is organized as follows: In Section 2, we construct a locally convex completion H of a finitely-generated $\mathbb{Z}$-graded vertex algebra $(V, Y, \mathbf{1})$ satisfying the standard grading-restriction axioms. In Section 3, we extend $Y(r_1^{L(0)}\cdot, z)r_2^{L(0)} : V \otimes V \to \overline{V}$ to continuous linear maps from $H\tilde{\otimes}H$ to H. In Section 4, we present the corresponding results for modules.

In this paper, we assume that the reader is familiar with the basic definitions and results in the theory of vertex algebras. The material in [B], [FLM] and [FHL] should be enough. We also assume that the reader is familiar with the basic definitions, constructions and results in the theory of locally convex topological vector spaces. The reader can find this material in, for example, [K1] and [K2].

We shall denote the set of integers, the set of real numbers and the set of complex numbers by the usual notations $\mathbb{Z}$, $\mathbb{R}$ and $\mathbb{C}$, respectively. We shall use i, j, k, l, m, n, p, q to denote integers. In particular, when we write, say, $k > 0$ (or $k \geq 0$), we mean that k is a positive integer (or a nonnegative integer). For a graded vector space V, we use V', V^* and $\overline{V}$ to denote the graded dual space, the dual space and the algebraic completion of V, respectively. For a topological vector space E, we use E^* to denote the topological dual space of E. The symbol $\otimes$ always denotes the vector space tensor product. The bifunctor given by completing the vector space tensor product of two topological vector spaces with the tensor product topology is denoted by $\tilde{\otimes}$.

Acknowledgment This research is supported in part by by NSF grant DMS-9622961.

1 A locally convex completion of a finitely-generated vertex algebra

In this section, we use "correlation functions" to construct the locally convex completion of a finitely-generated vertex algebra.

For any $k \geq 0$, let R_k be the space of rational functions in the complex variables $z_1, \dots, z_k$ with the only possible poles $z_i = z_j$ for $i \neq j$ and $z_i = 0, \infty$ $(i, j = 1, \dots, k)$. Let

$$M^k = \{(z_1, \dots, z_k) \in \mathbb{C}^k \mid z_i \neq z_j \text{ for } i \neq j; z_i \neq 0 \ (i, j = 1, \dots, k)\}$$

and let $\{K_n^{(k)}\}_{n>0}$, be a sequence of compact subsets of M^k satisfying

$$K_n^{(k)} \subset K_{n+1}^{(k)}, \quad n > 0,$$

and

$$M^k = \bigcup_{n>0} K_n^{(k)}.$$

For any $n > 0$, we define a map $\| \cdot \|_{R_k, n} : R_k \to [0, \infty)$ by

$$\|f\|_{R_k,n} = \max_{(z_1,\dots,z_k) \in K_n^{(k)}} |f(z_1, \dots, z_k)|$$

for $f \in R_k$. Then it is clear that $\| \cdot \|_{R_k, n}$ is a norm on R_k. Using this sequence of norms, we obtain a locally convex topology on R_k. Note that a sequence in R_k is convergent if and only if this sequence of functions is uniformly convergent on any compact subset of M^k. Clearly, this topology is independent of the choice of the sequence $\{K_n^{(k)}\}_{n>0}$.

Let V be a $\mathbb{Z}$-graded vertex algebra satisfying the standard grading-restriction axioms, that is,

$$V = \coprod_{n \in \mathbb{Z}} V_{(n)},$$

$$\dim V_{(n)} < \infty$$

for $n \in \mathbb{Z}$ and

$$V_{(n)} = 0$$

for n sufficiently small. By the duality properties of V, for any $v' \in V'$, any $u_1, \dots, u_k, v \in V$,

$$\langle v', Y(u_1, z_1) \cdots Y(u_k, z_k) v \rangle$$

is absolutely convergent in the region $|z_1| > \cdots > |z_k| > 0$ and can be analytically extended to an element

$$R(\langle v', Y(u_1, z_1) \cdots Y(u_k, z_k) v \rangle)$$

of R_k.

For any $u_1, \dots, u_k, v \in V$ and any $(z_1, \dots, z_k) \in M^k$, we have an element

$$Q(u_1, \dots, u_k, v; z_1, \dots, z_k) \in \overline{V}$$

defined by

$$\langle v', Q(z_1, \dots, z_k, v; z_1, \dots, z_k) \rangle = R(\langle v', Y(u_1, z_1) \cdots Y(u_k, z_k) v \rangle)$$

for $v' \in V'$. We denote the projections from V to $V_{(n)}$, $n \in +\mathbb{Z}$, by P_n. Let $\tilde{G}$ be the subspace of V^* consisting of linear functionals λ on V such that for any $k \geq 0$, $u_1, \dots, u_k, v \in V$,

$$\sum_{n \in \mathbb{Z}} \lambda(P_n(Q(u_1, \dots, u_k, v; z_1, \dots, z_k))) \tag{1.1}$$

is absolutely convergent for any $z_1, \dots, z_k$ in the region

$$M^k_{<1} = \{(z_1, \dots, z_k) \in M^k \mid |z_1|, \dots, |z_k| < 1\}.$$

The dual pair (V^*, V) of vector spaces gives V^* a locally convex topology. With the topology induced from the one on V^*, $\tilde{G}$ is also a locally convex space. Note that V' is a subspace of $\tilde{G}$.

For any $k \geq 0$, $\lambda \in \tilde{G}$, $u_1, \dots, u_k, v \in V$, both (1.1) and

$$\begin{aligned}\sum_{n\in\mathbb{Z}} \frac{\partial}{\partial z_1}\lambda(P_n(Q(u_1, \dots, u_k, v; z_1, \dots, z_k)))\\ = \sum_{n\in\mathbb{Z}} \lambda(P_n(Q(L(-1)u_1, \dots, u_k, v; z_1, \dots, z_k)))\end{aligned}$$

are absolutely convergent in the region $M^k_{<1}$. Thus (1.1) is analytic in z_1 when $(z_1, \dots, z_k)$ is in $M^k_{<1}$. Similarly, (1.1) is analytic in z_i for $i = 2, \dots, k$ when $(z_1, \dots, z_k)$ is in $M^k_{<1}$. So (1.1) defines an analytic function on $M^k_{<1}$ and we denote it by

$$g_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes v)$$

since (1.1) is multilinear in λ, $u_1, \dots, u_k$ and v. These functions span a vector space F_k of analytic functions on $M^k_{<1}$. So we obtain a linear map

$$g_k : \tilde{G} \otimes V^{\otimes(k+1)} \to F_k.$$

Fix a sequence $\{J^{(k)}_n\}_{n>0}$ of compact subsets of $M^k_{<1}$ such that

$$J^{(k)}_n \subset J^{(k)}_{n+1}, \quad n > 0,$$

and

$$\bigcup_{n>0} J^{(k)}_n = M^k_{<1}.$$

As in the case of R_k, using these compact subsets, we define a sequence of norms $\|\cdot\|_{F_k,n}$ on F_k, and these norms give a locally convex topology on F_k.

There is also an embedding ι_{F_k} from F_k to F_{k+1} defined as follows: We use $(z_0, \dots, z_k)$ instead of $(z_1, \dots, z_{k+1})$ to denote the elements of $M^{k+1}_{<1}$. For $\lambda \in \tilde{G}$, $u_1, \dots, u_k, v \in V$, since

$$Y(\mathbf{1}, z) = 1$$

for any nonzero complex number z,

$$g_{k+1}(\lambda \otimes \mathbf{1} \otimes u_1 \otimes \cdots \otimes u_k \otimes v)$$

as a function of $z_0, \dots, z_k$ is in fact independent of z_0, and is equal to

$$g_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes v)$$

as a function in $z_1, \dots, z_k$. Thus we obtain a well-defined linear map

$$\iota_{F_k} : F_k \to F_{k+1}$$

such that

$$\iota_{F_k} \circ g_k = g_{k+1} \circ \phi_k$$

where

$$\phi_k : \tilde{G} \otimes V^{\otimes(k+1)} \to \tilde{G} \otimes V^{\otimes(k+2)}$$

is defined by

$$\phi_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes v) = \lambda \otimes \mathbf{1} \otimes u_1 \otimes \cdots \otimes u_k \otimes v$$

for $\lambda \in \tilde{G}$, $u_1, \dots, u_k, v \in V$. It is clear that ι_{F_k} is injective. Thus we can regard F_k as a subspace of F_{k+1}. Moreover, we have:

Proposition 1.1 *For any $k \geq 0$, ι_{F_k} as a map from F_k to $\iota_{F_k}(F_k)$ is continuous and open. In other words, the topology on F_k is induced from that on F_{k+1}.*

Proof. We consider the two topologies on F_k, one is the topology defined above for F_k and the other induced from the topology on F_{k+1}. We need only prove that for any $n > 0$, (i) the norm $\|\cdot\|_{F_k,n}$ is continuous in the topology induced from the one on F_{k+1}, and (ii) the restriction of the norm $\|\cdot\|_{F_{k+1},n}$ to F_k is continuous in the topology on F_k.

Let $\{f_\alpha\}_{\alpha\in A}$ be a net in F_k convergent in the topology induced from the one on F_{k+1}. Then $\{f_\alpha\}_{\alpha\in A}$, when viewed as a net of functions in $z_0, z_1, \dots, z_k$, is convergent uniformly on any compact subset of M^{k+1}. Since f_α, $\alpha \in A$, are independent of z_0, $\{f_\alpha\}_{\alpha\in A}$ is in fact convergent uniformly on any compact subset of M^k, proving (i). Now let $\{f_\alpha\}_{\alpha\in A}$ be a net in F_k convergent in the topology on F_k. Then $\{f_\alpha\}_{\alpha\in A}$ is convergent uniformly on any compact subset of M^k. If we view f_α, $\alpha \in A$, as functions on $\mathbb{C} \times M^k$, then the net $\{f_\alpha\}_{\alpha\in A}$ is convergent uniformly on any subset of M^{k+1} of the form $\mathbb{C} \times K$ where K is a compact subset of M^k. But any compact subset of M^{k+1} is contained in a subset of the form $\mathbb{C} \times K$. So $\{f_\alpha\}_{\alpha\in A}$ is convergent uniformly on any compact subset of M^{k+1}, proving (ii). □

We equip the topological dual space F_k^*, $k \geq 0$, of F_k with the strong topology, that is, the topology of uniform convergence on all the weakly bounded subsets of F_k (see page 256 of [K1]). Then F_k^* is a locally convex space. In fact, F_k^* is a (DF)-space (see page 396 of [K1]).

For any $\lambda \in \tilde{G}$ and $u \in V$, let $Y_{-1}(u)\lambda \in \tilde{G}$ be defined by

$$(Y_{-1}(u)\lambda)(v) = \lambda((\mathrm{Res}_x x^{-1}Y(u, x))v)$$

for $v \in V$. (Note that $\mathrm{Res}_x x^{-1}Y(u, x)$ is u_{-1} in the usual notation. So we could denote $Y_{-1}(u)$ by u_{-1}. Here we use the notation $Y_{-1}(u)$ to avoid the possible confusion with the notations $u_0, u_1, \ldots$ for elements in V used later.) For $k \geq 0$, we define a linear map

$$\gamma_k : F_{k+1} \to F_k$$

by

$$\begin{aligned}\gamma_k(g_{k+1}(\lambda \otimes u_0 \otimes u_1 \otimes \cdots \otimes u_k \otimes v))\\ = g_k(Y_{-1}(u_0)\lambda) \otimes u_1 \otimes \cdots \otimes u_k \otimes v)\end{aligned}$$

for $\lambda \in \tilde{G}$, $u_0, u_1, \ldots, u_l, v \in V$.

Proposition 1.2 *The map γ_k is continuous and satisfies*

$$\gamma_k \circ \iota_{F_k} = I_{F_k} \tag{1.2}$$

where I_{F_k} is the identity map on F_k.

Proof. We use $(z_0, \ldots, z_k)$ instead of $(z_1, \ldots, z_{k+1})$ to denote the elements of $M^{k+1}_{<1}$. From the definition, we see that for any positive number $\epsilon < 1$, when $|z_1|, \ldots, |z_k| < \epsilon$,

$$\begin{aligned}&\gamma_k(g_{k+1}(\lambda \otimes u_0 \otimes u_1 \otimes \cdots \otimes u_l \otimes v))\\ &\quad = \frac{1}{2\pi\sqrt{-1}} \oint_{|z_0|=\epsilon} z_0^{-1} g_{k+1}(\lambda \otimes u_0 \otimes u_1 \otimes \cdots \otimes u_k \otimes v)dz_0\end{aligned}$$

for $\lambda \in \tilde{G}$, $u_0, \ldots, u_k, v \in V$. Thus for any $n > 0$, there exist $m_n > 0$ and positive number ϵ_n such that

$$\|\gamma_k(g_{k+1}(\lambda \otimes u_0 \otimes u_1 \otimes \cdots \otimes u_k \otimes v))\|_{F_k,n}$$

$$\begin{aligned}
&= \max_{(z_1,\dots,z_k)\in J_n^{(k)}} |\gamma_k(g_{k+1}(\lambda\otimes u_0\otimes u_1\otimes\cdots\otimes u_k\otimes v))| \\
&= \max_{(z_1,\dots,z_k)\in J_n^{(k)}} \left|\frac{1}{2\pi\sqrt{-1}}\oint_{|z_0|=\epsilon_n} z_0^{-1} g_{k+1}(\lambda\otimes u_0\otimes u_1\otimes\cdots\otimes u_k\otimes v)dz_0\right| \\
&\leq \max_{(z_1,\dots,z_k)\in J_n^{(k)},|z_0|=\epsilon_n} |g_{k+1}(\lambda\otimes u_0\otimes u_1\otimes\cdots\otimes u_k\otimes v)| \\
&\leq \max_{(z_0,\dots,z_k)\in J_{mn}^{(k+1)}} |g_{k+1}(\lambda\otimes u_0\otimes u_1\otimes\cdots\otimes u_k\otimes v)| \\
&= \|g_{k+1}(\lambda\otimes u_0\otimes u_1\otimes\cdots\otimes u_k\otimes v)\|_{F_{k+1},m_n}.
\end{aligned}$$

This inequality implies that γ_k is continuous.

For $\lambda\in\tilde{G}$, $u_1,\cdots,u_k,v\in V$, by definition,

$$\begin{aligned}
&g_{k+1}(\lambda\otimes\mathbf{1}\otimes u_1\otimes\cdots\otimes u_k\otimes v)\\
&\qquad=\iota_{F_k}(g_k(\lambda\otimes u_1\otimes\cdots\otimes u_k\otimes v)).
\end{aligned}$$

Since $Y_{-1}(\mathbf{1})\lambda=\lambda$,

$$\begin{aligned}
&\gamma_k(g_{k+1}(\lambda\otimes\mathbf{1}\otimes u_1\otimes\cdots\otimes u_k\otimes v))\\
&\qquad=g_k((Y_{-1}(\mathbf{1})\lambda)\otimes u_1\otimes\cdots\otimes u_k\otimes v)\\
&\qquad=g_k(\lambda\otimes u_1\otimes\cdots\otimes u_k\otimes v).
\end{aligned}$$

So we have (1.2). □

Corollary 1.3 *The adjoint map γ_k^* of γ_k satisfies*

$$\iota_{F_k}^*\circ\gamma_k^*=I_{F_k^*} \tag{1.3}$$

where

$$\iota_{F_k}^*:F_{k+1}^*\to F_k^*$$

is the adjoint of ι_{F_k} and $I_{F_k^*}$ is the identity on F_k^*. It is injective and continuous. As a map from F_k^* to $\gamma_k^*(F_k^*)$, it is also open. In particular, if we identify F_k^* with $\gamma_k^*(F_k^*)$, the topology on F_k^* is induced from the one on F_{k+1}^*.

Proof. The identity (1.3) is an immediate consequence of the identity (1.2). By this identity, we see that γ_k^* is injective. The continuity of γ_k^* is a consequence of the continuity of γ_k. To show that it is open as a map from F_k^* to $\gamma_k^*(F_k^*)$, we need only show that its inverse is continuous. Let $\{\mu_\alpha\}_{\alpha\in A}$

be a net in F_k^* such that $\{\gamma_k^*(\mu_\alpha)\}_{\alpha\in A}$ is convergent in F_{k+1}^*. Since $\iota_{F_k}^*$ is continuous, $\{\iota_{F_k}^*(\gamma_k^*(\mu_\alpha))\}_{\alpha\in A}$ is convergent in F_k^*. By the identity (1.2),

$$\mu_\alpha = \iota_{F_k}^*(\gamma_k^*(\mu_\alpha))$$

for $\alpha \in A$. Thus $\{\mu_\alpha\}_{\alpha\in A}$ is convergent in F_k^*, proving that the inverse of γ_k^* viewed as a map from F_k^* to $\gamma_k^*(F_k^*)$ is continuous. $\square$

We use $\langle \cdot, \cdot \rangle$ to denote the pairing between $\tilde{G}$ and the algebraic dual space $\tilde{G}^*$ of $\tilde{G}$. Since $V' \subset \tilde{G}$ and $V \subset \tilde{G}^*$, this pairing is an extension of the pairing between V' and V denoted using the same symbol. With this pairing, $\tilde{G}$ and $\tilde{G}^*$ form a dual pair of vector spaces. This dual pair of vector spaces gives a locally convex topology to $\tilde{G}^*$. Since $V' \subset \tilde{G}$, the dual space $\tilde{G}^*$ can be viewed as a subspace of $(V')^* = \overline{V}$. We define

$$e_k : V^{\otimes(k+1)} \otimes F_k^* \to \tilde{G}^* \subset \overline{V}$$

by

$$\langle \lambda, e_k(u_1 \otimes \cdots \otimes u_k \otimes v \otimes \mu)\rangle = \mu(g_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes v))$$

for $\lambda \in \tilde{G}$, $u_1, \dots, u_k, v \in V$ and $\mu \in F_k^*$.

We now assume that V is finitely generated. The generators of V span a finite-dimensional subspace X of V. We assume that the vacuum vector $\mathbf{1}$ is in X. Any norm on X induces a Banach space structure on X. Since X is finite-dimensional, all norms on X are equivalent, so that the topology induced by the norm is in fact independent of the norm.

Since X is a finite-dimensional Banach space and F_k^* is a locally convex space, $X^{\otimes(k+1)} \otimes F_k^*$ is also a locally convex space. We denote the image $e_k(X^{\otimes(k+1)} \otimes F_k^*)$ of $X^{\otimes(k+1)} \otimes F_k^* \subset V^{\otimes(k+1)} \otimes F_k^*$ under e_k by G_k.

Proposition 1.4 *For any $k \geq 0$, $G_k \subset G_{k+1}$.*

Proof. By definition,

$$\begin{aligned}
&\langle \lambda, e_k(u_1 \otimes \cdots \otimes u_k \otimes v \otimes \mu)\rangle \\
&\quad = \mu(g_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes v) \\
&\quad = \mu(\gamma_k(g_{k+1}(\lambda \otimes \mathbf{1} \otimes u_1 \otimes \cdots \otimes u_k \otimes v)) \\
&\quad = (\gamma_k^*(\mu))(g_{k+1}(\lambda \otimes \mathbf{1} \otimes u_1 \otimes \cdots \otimes u_k \otimes v)) \\
&\quad = \langle \lambda, e_k(\mathbf{1} \otimes u_1 \otimes \cdots \otimes u_k \otimes v \otimes \gamma_k^*(\mu))\rangle
\end{aligned}$$

for $\lambda \in \tilde{G}$, $u_1, \dots, u_k, v \in V$ and $\mu \in F_k^*$. Thus

$$e_k(u_1 \otimes \cdots \otimes u_k \otimes v \otimes \mu) = e_k(\mathbf{1} \otimes u_1 \otimes \cdots \otimes u_l \otimes v \otimes \gamma_k^*(\mu)) \in G_{k+1}$$

for $\lambda \in \tilde{G}$, $u_1, \dots, u_k, v \in V$ and $\mu \in F_k^*$. $\square$

Proposition 1.5 *The linear map*

$$e_k|_{X^{\otimes(k+1)} \otimes F_k^*} : X^{\otimes(k+1)} \otimes F_k^* \to \tilde{G}^*$$

is continuous.

Proof. By the definition of the locally convex topology on the tensor product $X^{\otimes(k+1)} \otimes F_k^*$, we need only prove that $e_k|_{X^{\otimes(k+1)} \otimes F_k^*}$ as a multilinear map from $X^{\otimes(k+1)} \times F_k^*$ to $\tilde{G}^*$ is continuous.

Let $\{(\mathcal{X}_\alpha, \mu_\alpha)\}_{\alpha \in A}$ be a net in $X^{\otimes(k+1)} \times F_k^*$ convergent to 0. Then the nets $\{\mathcal{X}_\alpha\}_{\alpha \in A}$ and $\{\mu_\alpha\}_{\alpha \in A}$ are convergent to 0 in $X^{\otimes(k+1)}$ and F_k^*, respectively. For any fixed $\lambda \in \tilde{G}$, since $X^{\otimes(k+1)}$ is a finite-dimensional Banach space, the linear map from $X^{\otimes(k+1)}$ to F_k defined by

$$u_1 \otimes \cdots \otimes u_k \otimes v \mapsto g_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes v)$$

for $u_1, \dots, u_k, v \in X$ is continuous. Thus $\{g_k(\lambda \otimes \mathcal{X}_\alpha)\}_{\alpha \in A}$ is convergent to 0 in F_k. In particular, there exists $\alpha_0 \in A$ such that $\{g_k(\lambda \otimes \mathcal{X}_\alpha)\}_{\alpha \in A, \alpha > \alpha_0}$ is weakly bounded. Thus

$$\left\{ \sup_{\alpha' \in A, \alpha' > \alpha_0} \mu_\alpha(g_k(\lambda \otimes \mathcal{X}_{\alpha'})) \right\}_{\alpha \in A}$$

is convergent to 0. In particular,

$$\{\mu_\alpha(g_k(\lambda \otimes \mathcal{X}_\alpha))\}_{\alpha \in A, \alpha > \alpha_0}$$

or equivalently

$$\{\mu_\alpha(g_k(\lambda \otimes \mathcal{X}_\alpha))\}_{\alpha \in A}$$

is convergent to 0. Since

$$\langle \lambda, e_k(\mathcal{X}_\alpha \otimes \mu_\alpha) \rangle = \mu_\alpha(g_k(\lambda \otimes \mathcal{X}_\alpha)),$$

we see that $\{\langle \lambda, e_k(\mathcal{X}_\alpha \otimes \mu_\alpha) \rangle\}_{\alpha \in A}$ is convergent to 0 for $\lambda \in \tilde{G}$. By the definition of the topology on $\tilde{G}^*$, $\{e_k(\mathcal{X}_\alpha \otimes \mu_\alpha)\}_{\alpha \in A}$ is convergent to 0 in $\tilde{G}^*$. $\square$

From Proposition 1.5, we conclude:

Corollary 1.6 *The quotient space*

$$(X^{\otimes(k+1)} \otimes F_k^*)/(e_k|_{X^{\otimes(k+1)}\otimes F_k^*})^{-1}(0)$$

is a locally convex space. $\square$

Since G_k is linearly isomorphic to

$$(X^{\otimes(k+1)} \otimes F_k^*)/(e_k|_{X^{\otimes(k+1)}\otimes F_k^*})^{-1}(0),$$

the locally convex space structure on

$$(X^{\otimes(k+1)} \otimes F_k^*)/(e_k|_{X^{\otimes(k+1)}\otimes F_k^*})^{-1}(0)$$

gives a locally convex space structure on G_k. Let H_k be the completion of G_k. Then H_k is a complete locally convex space.

Proposition 1.7 *The space H_k can be embedded canonically in H_{k+1}. The topology on H_k is the same as the one induced from the topology on H_{k+1}.*

Proof. The first conclusion follows from Proposition 1.4.

To prove the second conclusion, we need only prove that the topology on G_k is the same as the one induced from the topology on G_{k+1}. We have the following commutative diagram:

$$\begin{tikzcd}
X^{\otimes(k+1)} \otimes F_k^* \arrow[r, "\psi_k"] \arrow[d, "e_k"'] & X^{\otimes(k+2)} \otimes F_{k+1}^* \arrow[d, "e_{k+1}"] \\
G_k \arrow[r] & G_{k+1}
\end{tikzcd}$$

where

$$\psi_k : X^{\otimes(k+1)} \otimes F_k^* \to X^{\otimes(k+2)} \otimes F_{k+1}^*$$

is defined by

$$\psi_k(u_1 \otimes \cdots \otimes u_k \otimes v \otimes \mu) = \mathbf{1} \otimes u_1 \otimes \cdots \otimes u_k \otimes v \otimes \gamma_k^*(\mu)$$

for $u_1, \dots, u_k, v \in X$ and $\mu \in F_k^*$. By Corollary 1.3, if we identify F_k^* with $\gamma_k^*(F_k^*)$, the topology on F_k^* is induced from the one on F_{k+1}^*. By the definition of the topologies on G_k and G_{k+1} and the commutativity of the diagram above, we see that the topology on G_k is the same as the one induced from the topology on G_{k+1}. $\square$

By Proposition 1.7, we have a sequence $\{H_k\}_{k\geq 0}$ of strictly increasing complete locally convex spaces. Let

$$H = \bigcup_{k\geq 0} H_k.$$

We equip H with the inductive limit topology. Then H is a complete locally convex space. Let

$$G = \bigcup_{k\geq 0} G_k.$$

Then G is a dense subspace of H. Note that $V \subset G \subset \overline{V}$. Since elements of $\overline{V}$ are finite or infinite sums of elements of V, elements of G are also finite or infinite sums of elements of V. Thus infinite sums of elements of V belonging to G must be convergent in the topology on H. So G is in the closure of V. Since G is dense in H, we obtain:

Theorem 1.8 *The vector space H equipped with the strict inductive limit topology is a locally convex completion of V.* $\square$

2 The locally convex completion and the vertex operator map

In this section we construct continuous linear maps from the topological completion of $H \otimes H$ to H associated with conformal equivalence classes of closed disks with two ordered open disks inside deleted and with the standard parametrizations at the boundary components.

We consider the closed unit disk centered at 0. We delete two ordered open disks inside: the first centered at z with radius r_1 and the second centered at 0 with radius r_2. See the figure below.



The positive numbers r_1, r_2 and the nonzero complex number z must satisfy the conditions $r_2 + 2r_1 < 1$ and $r_2 < |z| < 1$. The three boundary circles are parametrized by the maps

$$\begin{array}{rcl} e^{i\theta} & \mapsto & e^{i\theta} \\ e^{i\theta} & \mapsto & z + r_1 e^{i\theta} \\ e^{i\theta} & \mapsto & r_2 e^{i\theta}. \end{array}$$

We denote the resulting closed disk with two disjoint open disks inside deleted and with ordered boundary components parametrized as above by $D(z; r_1, r_2)$. Note that any closed disks with two ordered open disks inside deleted and with the standard parametrizations is conformally equivalent to $D(z; r_1, r_2)$ for some $z \in \mathbb{C}^\times$, and some positive numbers r_1, r_2 satisfying $r_2 + 2r_1 < 1$ and $r_2 < |z| < 1$.

Let $H \tilde{\otimes} H$ be the topological completion of the vector space tensor product $H \otimes H$. We would like to construct a continuous linear map

$$\overline{\nu}_Y([D(z; r_1, r_2)]) : H \tilde{\otimes} H \to H$$

associated with the conformal equivalence class $[D(z; r_1, r_2)]$ of $D(z; r_1, r_2)$. We know that $D(z; r_1, r_2)$ corresponds to the sphere $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ with the negatively oriented puncture ∞, the ordered positively oriented puncture $z, 0$

and with the local coordinates $w \mapsto 1/w$, $w \mapsto (w-z)/r_1$ and w/r_2 vanishing at ∞, z and 0, respectively. In the notation of [H6], the conformal equivalence class of this sphere with tubes of type $(1,2)$ is $(z;\mathbf{0},(1/r_1,\mathbf{0}),(1/r_2,\mathbf{0})) \in K(2)$. Associated with this class, we have a linear map

$$\nu_Y((z;\mathbf{0},(1/r_1,\mathbf{0}),(1/r_2,\mathbf{0}))) = Y(r_1^{L(0)}\cdot, z)r_2^{L(0)}\cdot : V\otimes V \to \overline{V}$$

(see [H6]). We now show that this linear map can be extended to $H\tilde{\otimes}H$, that the image of this extension is in H and that this extension is continuous. Then we define $\overline{\nu}_Y([D(z;r_1,r_2)])$ to be this extension.

For any $\lambda \in \tilde{G}$ and $u \in V$, we define an element $u \diamond_{[D(z;r_1,r_2)]} \lambda \in V^*$ by

$$\begin{aligned}(u \diamond_{[D(z;r_1,r_2)]} \lambda)(v) &= \sum_{n\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u, z)r_2^{L(0)}v))\\ &= \sum_{n\in\mathbb{Z}} \lambda(P_n(Q(r_1^{L(0)}u, r_2^{L(0)}v; z)))\end{aligned}$$

for $v \in V$.

Proposition 2.1 *The element* $u \diamond_{[D(z;r_1,r_2)]} \lambda$ *is in* $\tilde{G}$.

Proof. For any $k \geq 0$, $u_1, \dots, u_k, v \in V$,

$$\begin{aligned}&\sum_{m\in\mathbb{Z}} (u \diamond_{[D(z;r_1,r_2)]} \lambda)(P_m(Q(u_1,\dots,u_k,v;z_1,\dots,z_k)))\\ &\quad= \sum_{m\in\mathbb{Z}}\sum_{n\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u,z)r_2^{L(0)}P_m(Q(u_1,\dots,u_k,v;z_1,\dots,z_k)))\\ &\quad= \sum_{m\in\mathbb{Z}}\sum_{n\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u,z)\cdot\\ &\qquad\qquad \cdot P_m(Q(r_2^{L(0)}u_1,\dots,r_2^{L(0)}u_k,r_2^{L(0)}v;r_2z_1,\dots,r_2z_k))). \quad (2.1)\end{aligned}$$

Since the $\mathbb{Z}$-grading of V is lower-truncated and since $r_2 < |z| < 1$, for any $(z_1,\dots,z_k) \in M^k_{<1}$, there exists a positive number $\delta_2 > 1$ such that the Laurent series in t_2

$$\begin{aligned}&\sum_{m\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u,z)P_m(Q(r_2^{L(0)}u_1,\dots,r_2^{L(0)}u_k,r_2^{L(0)}v;r_2z_1,\dots,r_2z_k))))t_2^m\\ &\quad= \sum_{m\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u,z)\cdot\\ &\qquad\qquad \cdot P_m(t_2^{L(0)}Q(r_2^{L(0)}u_1,\dots,r_2^{L(0)}u_k,r_2^{L(0)}v;r_2z_1,\dots,r_2z_k))))\\ &\quad= \sum_{m\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u,z)P_m(Q((t_2r_2)^{L(0)}u_1,\dots,(t_2r_2)^{L(0)}u_k,(t_2r_2)^{L(0)}v;\\ &\qquad\qquad t_2r_2z_1,\dots,t_2r_2z_k))))\end{aligned}$$

in t_2 has only finitely many negative powers and is absolutely convergent to $\lambda(P_n(Q(r_1^{L(0)}u, (t_2r_2)^{L(0)}u_1, \ldots, (t_2r_2)^{L(0)}u_k, (t_2r_2)^{L(0)}v; z, t_2r_2z_1, \ldots, t_2r_2z_k)))$ when $0 < |t_2| < \delta_2$. Since $\lambda \in \tilde{G}$ and $|z| < 1$, for any $(z_1, \ldots, z_k) \in M^k_{<1}$, there exists a positive number $\delta_1 > 1$ such that the Laurent series in t_1

$$\begin{aligned}
&\sum_{n\in\mathbb{Z}} \lambda(P_n(Q(r_1^{L(0)}u, r_2^{L(0)}u_1, \ldots, r_2^{L(0)}u_k, r_2^{L(0)}v; z, r_2z_1, \ldots, r_2z_k)))t_1^n\\
&\quad = \sum_{n\in\mathbb{Z}} \lambda(P_n(t_1^{L(0)}Q(r_1^{L(0)}u, r_2^{L(0)}u_1, \ldots, r_2^{L(0)}u_k, r_2^{L(0)}v; z, r_2z_1, \ldots, r_2z_k)))\\
&\quad = \sum_{n\in\mathbb{Z}} \lambda(P_n(Q((t_1r_1)^{L(0)}u, (t_1r_2)^{L(0)}u_1, \ldots, (t_1r_2)^{L(0)}u_k, (t_1r_2)^{L(0)}v;\\
&\qquad\qquad t_1z, t_1r_2z_1, \ldots, t_1r_2z_k)))
\end{aligned}$$

is absolutely convergent when $0 < |t_1| < \delta_1$. Thus for $(z_1, \ldots, z_k) \in M^k_{<1}$, the iterated sum

$$\begin{aligned}
\sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u, z)P_m(Q(r_2^{L(0)}u_1, \ldots, r_2^{L(0)}u_k, r_2^{L(0)}v;\\
r_2z_1, \ldots, r_2z_k))))t_1^nt_2^m
\end{aligned} \tag{2.2}$$

is absolutely convergent when $0 < |t_1| < \delta_1$ and $0 < |t_2| < \delta_2$.

The iterated sum of (2.2) gives a function of t_1 and t_2 in the region $0 < |t_1| < \delta_1$, $0 < |t_2| < \delta_2$. It is clear that this function is analytic in t_1 and t_2. Thus it has a Laurent expansion which must be

$$\begin{aligned}
\sum_{n,m\in\mathbb{Z}} \lambda(P_n(Y(r_1^{L(0)}u, z)P_m(Q(r_2^{L(0)}u_1, \ldots, r_2^{L(0)}u_k, r_2^{L(0)}v;\\
r_2z_1, \ldots, r_2z_k))))t_1^nt_2^m.
\end{aligned}$$

Since this double sum is equal to the Laurent expansion, it is absolutely convergent and thus both iterated sums are absolutely convergent. In particular, when $t_1 = t_2 = 1$, we see that the right-hand side of (2.1) and consequently the left-hand side of (2.1) are absolutely convergent when $(z_1, \ldots, z_k) \in M^k_{<1}$, proving that $u \diamond_{[D(z;r_1,r_2)]} \lambda$ is in $\tilde{G}$. $\square$

For any $l \geq 0$, we define a linear map $\alpha_l : \tilde{G} \otimes X^{l+1} \otimes F_l^* \to V^*$ by

$$\begin{aligned}
&(\alpha_l(\lambda \otimes v_1 \otimes \ldots \otimes v_l \otimes v \otimes \nu))(u)\\
&\qquad = \langle u \diamond_{[D(z;r_1,r_2)]} \lambda, e_l(v_1 \otimes \ldots \otimes v_l \otimes v \otimes \nu)\rangle.
\end{aligned}$$

for $\lambda \in \tilde{G}$, $v_1, \ldots, v_l, v \in X$, $\nu \in F_l^*$ and $u \in V$.

Proposition 2.2 *The image of α_l is in $\tilde{G}$.*

Proof. For any $k \geq 0$, $\lambda \in \tilde{G}$, $u_1, \ldots, u_k, u, \in V$, $v_1, \ldots, v_l, v \in X$ and $\nu \in F_l^*$,

$$\begin{aligned}
&\sum_{n\in\mathbb{Z}}(\alpha_l(\lambda \otimes v_1 \otimes \ldots \otimes v_l \otimes v \otimes \nu))(P_n(Q(u_1, \ldots, u_k, u; z_1, \ldots, z_k)))\\
&\quad= \sum_{n\in\mathbb{Z}}\langle(P_n(Q(u_1, \ldots, u_k, u; z_1, \ldots, z_k)) \diamond_{[D(z;r_1,r_2)]} \lambda),\\
&\qquad\qquad\qquad e_l(v_1 \otimes \cdots \otimes v_l \otimes v \otimes \nu)\rangle\\
&\quad= \sum_{n\in\mathbb{Z}}\nu(g_l((P_n(Q(u_1, \ldots, u_k, u; z_1, \ldots, z_k)) \diamond_{[D(z;r_1,r_2)]} \lambda)\\
&\qquad\qquad\qquad \otimes v_1 \otimes \cdots \otimes \otimes v_l \otimes v))\\
&\quad= \sum_{n\in\mathbb{Z}}\nu\left(\sum_{m\in\mathbb{Z}}(P_n(Q(u_1, \ldots, u_k, u; z_1, \ldots, z_k)) \diamond_{[D(z;r_1,r_2)]} \lambda)\right.\\
&\qquad\qquad\qquad \left.(P_m(Q(v_1, \ldots, v_k, v; z_{k+1}, \ldots, z_{k+l})))\right). \qquad (2.3)
\end{aligned}$$

By the definition of

$$P_n(Q(u_1, \ldots, u_k, u; z_1, \ldots, z_k)) \diamond_{[D(z;r_1,r_2)]} \lambda,$$

we have

$$\begin{aligned}
&(P_n(Q(u_1, \ldots, u_k, u; z_1, \ldots, z_k)) \diamond_{[D(z;r_1,r_2)]} \lambda)\\
&\qquad\qquad (P_m(Q(v_1, \ldots, v_k, v; z_{k+1}, \ldots, z_{k+l})))\\
&\quad= \sum_{p\in\mathbb{Z}}\lambda(P_p(Y(r_1^{L(0)}P_n(Q(u_1, \ldots, u_k, u; z_1, \ldots, z_k)), z) \cdot\\
&\qquad\qquad \cdot r_2^{L(0)}P_m(Q(v_1, \ldots, v_l, v; z_{k+1}, \ldots, z_{k+l}))))\\
&\quad= \sum_{p\in\mathbb{Z}}\lambda(P_p(Y(P_n(Q(r_1^{L(0)}u_1, \ldots, r_1^{L(0)}u_k, r_1^{L(0)}u; r_1z_1, \ldots, r_1z_k)), z) \cdot\\
&\qquad\qquad \cdot P_m(Q(r_2^{L(0)}v_1, \ldots, r_2^{L(0)}v_l, r_2^{L(0)}v; r_2z_{k+1}, \ldots, r_2z_{k+l})))).
\end{aligned} \qquad (2.4)$$

Using the associativity of vertex operators, we know that the element

$$\begin{aligned}
&\sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}}Y(P_n(Q(r_1^{L(0)}u_1, \ldots, r_1^{L(0)}u_k, r_1^{L(0)}u; r_1z_1, \ldots, r_1z_k)), z)\cdot\\
&\qquad\qquad \cdot P_m(Q(r_2^{L(0)}v_1, \ldots, r_2^{L(0)}v_l, r_2^{L(0)}v; r_2z_{k+1}, \ldots, r_2z_{k+l}))
\end{aligned}$$

in $\overline{V}$ is equal to

$$Q(r_1^{L(0)}u_1, \dots, r_1^{L(0)}u_k, r_1^{L(0)}u, r_2^{L(0)}v_1, \dots, r_2^{L(0)}v_l, r_2^{L(0)}v; \\ r_1z_1 + z, \dots, r_1z_k + z, z, r_2z_{k+1}, \dots, r_2z_{k+l})$$

when $(z_1, \dots, z_k) \in M^k_{<1}$ and $(z_{k+1}, \dots, z_{k+l}) \in M^l_{<1}$. This fact and (2.4) imply that

$$\sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}}(P_n(Q(u_1, \dots, u_k, u; z_1, \dots, z_k)) \diamond_{[D(z;r_1,r_2)]} \lambda) \\ (P_m(Q(v_1, \dots, v_k, v; z_{k+1}, \dots, z_{k+l})))$$

is convergent absolutely to

$$\lambda(P_p(Q(r_1^{L(0)}u_1, \dots, r_1^{L(0)}u_k, r_1^{L(0)}u, r_2^{L(0)}v_1, \dots, r_2^{L(0)}v_l, r_2^{L(0)}v; \\ r_1z_1 + z, \dots, r_1z_k + z, z, r_2z_{k+1}, \dots, r_2z_{k+l})))$$

when $(z_1, \dots, z_k) \in M^k_{<1}$ and $(z_{k+1}, \dots, z_{k+l}) \in M^l_{<1}$. Since $\lambda \in \tilde{G}$,

$$\sum_{p\in\mathbb{Z}}\lambda(P_p(Q(r_1^{L(0)}u_1, \dots, r_1^{L(0)}u_k, r_1^{L(0)}u, r_2^{L(0)}v_1, \dots, r_2^{L(0)}v_l, r_2^{L(0)}v; \\ r_1z_1 + z, \dots, r_1z_k + z, z, r_2z_{k+1}, \dots, r_2z_{k+l})))$$

is absolutely convergent when $(z_1, \dots, z_k) \in M^k_{<1}$ and $(z_{k+1}, \dots, z_{k+l}) \in M^l_{<1}$. The arguments above prove that the iterated sum

$$\sum_{p\in\mathbb{Z}}\sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}}\lambda(P_p(Y(P_n(Q(r_1^{L(0)}u_1, \dots, r_1^{L(0)}u_k, r_1^{L(0)}u; r_1z_1, \dots, r_1z_k)), z)\cdot \\ \cdot P_m(Q(r_2^{L(0)}v_1, \dots, r_2^{L(0)}v_l, r_2^{L(0)}v; r_2z_{k+1}, \dots, r_2z_{k+l}))))$$

is absolutely convergent when $(z_1, \dots, z_k) \in M^k_{<1}$ and $(z_{k+1}, \dots, z_{k+l}) \in M^l_{<1}$. The same method as in the proof of Proposition 2.1 shows that in fact the corresponding triple sum is absolutely convergent and thus all the iterated sums are absolutely convergent and are all equal when $(z_1, \dots, z_k) \in M^k_{<1}$ and $(z_{k+1}, \dots, z_{k+l}) \in M^l_{<1}$. So

$$\sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}}\sum_{p\in\mathbb{Z}}\lambda(P_p(Y(P_n(Q(r_1^{L(0)}u_1, \dots, r_1^{L(0)}u_k, r_1^{L(0)}u; r_1z_1, \dots, r_1z_k)), z)\cdot \\ \cdot P_m(Q(r_2^{L(0)}v_1, \dots, r_2^{L(0)}v_l, r_2^{L(0)}v; r_2z_{k+1}, \dots, r_2z_{k+l})))) \\ = \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}}(P_n(Q(u_1, \dots, u_k, u; z_1, \dots, z_k)) \diamond_{[D(z;r_1,r_2)]} \lambda) \\ (P_m(Q(v_1, \dots, v_k, v; z_{k+1}, \dots, z_{k+l})))$$

is absolutely convergent when $(z_1, \dots, z_k) \in M^k_{<1}$ and $(z_{k+1}, \dots, z_{k+l}) \in M^l_{<1}$. By (2.3), we see that the left-hand side of (2.3) is also absolutely convergent when $(z_1, \dots, z_k) \in M^k_{<1}$ and $(z_{k+1}, \dots, z_{k+l}) \in M^l_{<1}$, proving that

$$\alpha_l(\lambda \otimes v_1 \otimes \ldots \otimes v_l \otimes v \otimes \nu)$$

is indeed in $\tilde{G}$. $\square$

By Proposition 2.2,

$$\sum_{n\in\mathbb{Z}} \alpha_l(\lambda \otimes v_1 \otimes \ldots \otimes v_l \otimes v \otimes \nu)(P_n(Q(u_1, \dots, u_k, u; z_1, \dots, z_k)))$$

is absolutely convergent and the sum is equal to

$$g_k(\alpha_l(\lambda \otimes v_1 \otimes \ldots \otimes v_l \otimes v \otimes \nu) \otimes u_1 \otimes \cdots \otimes u_k \otimes u) \in F_k.$$

We define an linear map

$$\beta_{k,l} : F^*_k \otimes F^*_l \to F^*_{k+l+1}$$

by

$$\begin{aligned}(\beta_{k,l}(\mu, \nu))(g_{k+l+1}(\lambda \otimes u_1 \otimes \ldots \otimes u_{k+1} \otimes u \otimes v_1 \otimes \cdots \otimes v_l \otimes v))\\ = \mu(g_k(\alpha_l(\lambda \otimes v_1 \otimes \ldots \otimes v_l \otimes v \otimes \nu) \otimes u_1 \otimes \cdots \otimes u_k \otimes u))\end{aligned}$$

for $\lambda \in \tilde{G}$, $u_1, \dots, u_k, u, v_1, \dots, v_l, v \in V$, $\mu \in F^*_k$ and $\nu \in F^*_l$. In fact this formula only gives a linear map from $F^*_k \otimes F^*_l$ to the algebraic dual of F_{k+l+1}. We have:

Proposition 2.3 *The image of the map $\beta_{k,l}$ is indeed in F^*_{k+l+1} and the map $\beta_{k,l}$ is continuous.*

Proof. To avoid notational confusions in the proof, we use F^ζ_k, $k \geq 0$, to denote the space F_k whose elements are viewed as functions of $(\zeta_1, \dots, \zeta_k) \in M^k_{<1}$. Similarly we have the notation F^η_l, $l \geq 0$. Let $\mu \in F^*_k$ and $\nu \in F^*_l$. For any $\lambda \in \tilde{G}$, $u_1, \dots, u_k, u$, $v_1, \dots, v_l, v \in V$, we have an element

$$g(z_1, \dots, z_{k+l+1}) = g_{k+l+1}(\lambda \otimes u_1 \otimes \ldots \otimes u_{k+1} \otimes u \otimes v_1 \otimes \cdots \otimes v_l \otimes v) \quad (2.5)$$

of F_{k+l+1}. By definition

$$\begin{aligned}
&|(\beta_{k,l}(\mu,\nu))(g(z_1,\dots,z_{k+l+1}))|\\
&\quad=|(\beta_{k,l}(\mu,\nu))(g_{k+l+1}(\lambda\otimes u_1\otimes\ldots\otimes u_{k+1}\otimes u\otimes v_1\otimes\cdots\otimes v_l\otimes v))|\\
&\quad=|\mu(g_k(\alpha_l(\lambda\otimes v_1\otimes\ldots\otimes v_l\otimes v\otimes\nu)\otimes u_1\otimes\cdots\otimes u_k\otimes u))|. \qquad (2.6)
\end{aligned}$$

We now view

$$g_k(\alpha_l(\lambda\otimes v_1\otimes\ldots\otimes v_l\otimes v\otimes\nu)\otimes u_1\otimes\cdots\otimes u_k\otimes u)$$

as an element of F_k^ζ. Then

$$\begin{aligned}
&g_k(\alpha_l(\lambda\otimes v_1\otimes\ldots\otimes v_l\otimes v\otimes\nu)\otimes u_1\otimes\cdots\otimes u_k\otimes u)\\
&\quad=\sum_{p\in\mathbb{Z}}(\alpha_l(\lambda\otimes v_1\otimes\ldots\otimes v_l\otimes v\otimes\nu))(P_p(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_k)))\\
&\quad=\sum_{p\in\mathbb{Z}}\nu(g_l((P_p(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_k))\diamond_{[D(z;r_1,r_2)]}\lambda)\\
&\qquad\qquad\otimes v_1\otimes\cdots\otimes v_l\otimes v)). \qquad (2.7)
\end{aligned}$$

For fixed $\zeta_1,\dots,\zeta_k$, we view

$$g_l((P_p(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_{k_1}))\diamond_{[D(z;r_1,r_2)]}\lambda)\otimes v_1\otimes\cdots\otimes v_l\otimes v)$$

as an element of F_l^η. Then by definition we have

$$\begin{aligned}
&\sum_{p\in\mathbb{Z}}g_l((P_p(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_{k_1}))\diamond_{[D(z;r_1,r_2)]}\lambda)\otimes v_1\otimes\cdots\otimes v_l\otimes v)\\
&\quad=\sum_{p\in\mathbb{Z}}\sum_{q\in\mathbb{Z}}(P_p(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_{k_1}))\diamond_{[D(z;r_1,r_2)]}\lambda)\\
&\qquad\qquad(P_q(Q(v_1,\dots,v_l,v;\eta_1,\dots,\eta_l)))\\
&\quad=\sum_{p\in\mathbb{Z}}\sum_{q\in\mathbb{Z}}\sum_{j\in\mathbb{Z}}\lambda(P_j(Y(r_1^{L(0)}P_p(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_{k_1})),z)\cdot\\
&\qquad\qquad\cdot r_2^{L(0)}P_q(Q(v_1,\dots,v_l,v;\eta_1,\dots,\eta_l))))\\
&\quad=\sum_{p\in\mathbb{Z}}\sum_{q\in\mathbb{Z}}\sum_{j\in\mathbb{Z}}\lambda(P_j(Y(P_p(Q(r_1^{L(0)}u_1,\dots,r_1^{L(0)}u_k,r_1^{L(0)}u;r_1\zeta_1,\dots,r_1\zeta_{k_1})),z)\cdot\\
&\qquad\qquad\cdot P_q(Q(r_2^{L(0)}v_1,\dots,r_2^{L(0)}v_l,r_2^{L(0)}v;r_2\eta_1,\dots,r_2\eta_l)))). \qquad (2.8)
\end{aligned}$$

We now calculate the right-hand side of (2.8). For any nonzero complex numbers t_0,t_1,t_2,

$$\sum_{p,q,j\in\mathbb{Z}}\lambda(P_j(Y(P_p(Q(r_1^{L(0)}u_1,\dots,r_1^{L(0)}u_k,r_1^{L(0)}u;r_1\zeta_1,\dots,r_1\zeta_{k_1})),z)\cdot$$

$$
\begin{aligned}
&\qquad\qquad \cdot P_q(Q(r_2^{L(0)}v_1, \ldots, r_2^{L(0)}v_l, r_2^{L(0)}v; r_2\eta_1, \ldots, r_2\eta_l))))t_0^j t_1^p t_2^q\\
&= \sum_{p,q,j\in\mathbb{Z}} \lambda(P_j(t_0^{L(0)}Y(P_p(t_1^{L(0)}Q(r_1^{L(0)}u_1, \ldots, r_1^{L(0)}u_k, r_1^{L(0)}u;\\
&\qquad\qquad r_1\zeta_1, \ldots, r_1\zeta_{k_1})), z) \cdot\\
&\qquad\qquad \cdot P_q(t_2^{L(0)}Q(r_2^{L(0)}v_1, \ldots, r_2^{L(0)}v_l, r_2^{L(0)}v; r_2\eta_1, \ldots, r_2\eta_l))))\\
&= \sum_{p,q,j\in\mathbb{Z}} \lambda(P_j(Y(P_p(Q((t_0t_1r_1)^{L(0)}u_1, \ldots, (t_0t_1r_1)^{L(0)}u_k, (t_0t_1r_1)^{L(0)}u;\\
&\qquad\qquad t_0t_1r_1\zeta_1, \ldots, t_0t_1r_1\zeta_{k_1})), t_0z) \cdot\\
&\qquad\qquad \cdot P_q(Q((t_0t_2r_1)^{L(0)}v_1, \ldots, (t_0t_2r_1)^{L(0)}v_l, (t_0t_2r_1)^{L(0)}v;\\
&\qquad\qquad t_0t_2r_2\eta_1, \ldots, t_0t_2r_2\eta_l)))). \qquad (2.9)
\end{aligned}
$$

By the associativity of vertex operators and the definition of the map g_{k+l+1}, there exists real numbers $\delta_0, \delta_1, \delta_2 > 1$ such that the right-hand side of (2.9) is convergent absolutely to

$$
\begin{aligned}
&g_{k+l+1}(\lambda \otimes (t_0t_1r_1)^{L(0)}u_1 \otimes \cdots \otimes (t_0t_1r_1)^{L(0)}u_k \otimes (t_0t_1r_1)^{L(0)}u \otimes\\
&\qquad (t_0t_2r_1)^{L(0)}v_1 \otimes \cdots \otimes (t_0t_2r_1)^{L(0)}v_l \otimes\\
&\qquad (t_0t_2r_1)^{L(0)}v)\big|_{z_i=t_0z+t_0t_1r_1\zeta_i, i=1,\ldots,k, z_{k+1}=t_0z, z_{k+1+j}=t_0t_2r_2\eta_j, j=1,\ldots,l}
\end{aligned}
\tag{2.10}
$$

when $0 < t_0 < \delta_0$, $0 < t_1 < \delta_1$ and $0 < t_1 < \delta_1$. In particular, when $t_0 = t_1 = t_2 = 1$, we see that (2.10) is equal to and thus the right-hand side of (2.8) is convergent absolutely to

$$
\begin{aligned}
&g_{k+l+1}(\lambda \otimes r_1^{L(0)}u_1 \otimes \cdots \otimes r_1^{L(0)}u_k \otimes r_1^{L(0)}u \otimes r_2^{L(0)}v_1 \otimes \cdots\\
&\qquad \otimes r_2^{L(0)}v_l \otimes r_2^{L(0)}v)\big|_{z_i=z+r_1\zeta_i, i=1,\ldots,k, z_{k+1}=z, z_{k+1+j}=r_2\eta_j, j=1,\ldots,l}.
\end{aligned}
\tag{2.11}
$$

Hence for fixed $(\zeta_1, \ldots, \zeta_k) \in M^k_{<1}$, (2.11) as a function of $\eta_1, \ldots, \eta_l$ is an element of F^η_l. We denote this element by $f_{g;\zeta_1,\ldots,\zeta_k}(\eta_1, \ldots, \eta_l)$.

We now view μ and ν as elements of $(F^\zeta_k)^*$ and $(F^\eta_l)^*$, respectively. Since ν is continuous, by the calculations above, we have

$$
\begin{aligned}
&\nu(f_{g;\zeta_1,\ldots,\zeta_k}(\eta_1, \ldots, \eta_l))\\
&\qquad = \nu\Bigg(\sum_{p\in\mathbb{Z}} g_l((P_p(Q(u_1, \ldots, u_k, u; \zeta_1, \ldots, \zeta_{k_1})) \diamond_{[D(z;r_1,r_2)]} \lambda)
\end{aligned}
$$

$$
\begin{aligned}
&\qquad\qquad \otimes v_1 \otimes \cdots \otimes v_l \otimes v)\Big)\\
&= \sum_{p\in\mathbb{Z}} \nu(g_l((P_p(Q(u_1, \dots, u_k, u; \zeta_1, \dots, \zeta_{k_1})) \diamond_{[D(z;r_1,r_2)]} \lambda)\\
&\qquad\qquad \otimes v_1 \otimes \cdots \otimes v_l \otimes v)). \qquad (2.12)
\end{aligned}
$$

By the calculations from (2.6) to (2.12), we see that

$$\nu(f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l)) \in F_k^{\zeta}$$

and we obtain

$$
\begin{aligned}
&|(\beta_{k,l}(\mu, \nu))(g(z_1, \dots, z_{k+l+1}))|\\
&\quad = |\mu(\nu(f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l)))|. \qquad (2.13)
\end{aligned}
$$

For $g(z_1, \dots, z_{k+l+1}) \in F_{k+l+1}$ not of the form (2.5), we use linearity to define the functions $f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l)$. Then (2.13) holds for any $g(z_1, \dots, z_{k+l+1}) \in F_{k+l+1}$.

From the definition of $f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l)$, we see that for any fixed $(\zeta_1, \dots, \zeta_k) \in M^k_{<1}$, the linear map from F_{k+l+1} to F_l^{η} given by

$$g \mapsto f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l)$$

for $g \in F_{k+l+1}$ is continuous, and for any fixed $\nu \in (F_l^{\eta})^*$, the linear map from F_{k+l+1} to F_k^{ζ} given by

$$g \mapsto \nu(f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l))$$

is also continuous. Thus by (2.13), $\beta_{k,l}(\mu, \nu)$ is continuous. So the elements of image of $\beta_{k,l}$ are in F^*_{k+l+1}.

From the definition of $f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l)$, we see that given any weakly bounded subset B of F_{k+l+1}, the subset

$$B' = \{f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l) \mid g \in B\} \subset F_l^{\eta}$$

for fixed $(\zeta_1, \dots, \zeta_k) \in M^k_{<1}$ is weakly bounded, and thus for any net $\{\nu_\alpha\}_{\alpha\in A}$ convergent to 0 in $(F_l^{\eta})^*$, there exists $\alpha_0 \in A$ such that the subset

$$B''(\{\nu_\alpha\}_{\alpha\in A}) = \{\nu_\alpha(f_{g;\zeta_1,\dots,\zeta_k}(\eta_1, \dots, \eta_l)) \mid g \in B, \alpha > \alpha_0\} \subset F_k^{\zeta}$$

is also weakly bounded.

Now let $\{(\mu_\alpha, \nu_\alpha)\}_{\alpha\in A}$ be a net convergent to 0 in $(F_k^\zeta)^* \times (F_l^\eta)^*$. Then the nets $\{\mu_\alpha\}_{\alpha\in A}$ and $\{\nu_\alpha\}_{\alpha\in A}$ are convergent to 0 in (F_k^ζ) and (F_l^η), respectively. Thus by (2.13) and the discussion above,

$$\begin{aligned} &\sup\nolimits_{g(z_1,\dots,z_{k+l+1})\in B} |(\beta_{k,l}(\mu_\alpha, \nu_\alpha))(g(z_1,\dots,z_{k+l+1}))| \\ &\quad\le \sup\nolimits_{h(\zeta_1,\dots,\zeta_k)\in B''(\{\nu_\alpha\}_{\alpha\in A})} |\mu_\alpha(h(\zeta_1,\dots,\zeta_k))| \end{aligned} \tag{2.14}$$

when $\alpha > \alpha_0$. By the definition of the topology on F_k^*, the net on the right-hand side of (2.14) is convergent to 0. Thus the net on the left-hand side of (2.14) is convergent to 0, proving the continuity of the map β_{k_1,k_2}. $\square$

Let

$$h_1 = e_k(u_1 \otimes \cdots \otimes u_k \otimes u \otimes \mu) \in G_k$$

and

$$h_2 = e_l(v_1 \otimes \cdots \otimes v_l \otimes v \otimes \nu) \in G_l$$

where $u_1, \dots, u_k, u, v_1, \dots, v_l, v \in X$, $\mu \in F_k^*$ and $\mu \in F_l^*$. We define

$$\begin{aligned} &(\overline{\nu}_Y([D(z; r_0, r_1, r_2)]))(h_1 \otimes h_2) \\ &\quad = e_{k+l+1}(u_1 \otimes \cdots \otimes u_k \otimes u \otimes v_1 \otimes \cdots \otimes v_l \otimes v \otimes \beta_{k,l}(\mu, \nu)). \end{aligned}$$

Note that any element of G_k or G_l is a linear combination of elements of the form h_1 or h_2, respectively, given above, and that k and l are arbitrary. Thus we obtain a linear map

$$\overline{\nu}_Y([D(z; r_0, r_1, r_2)])\big|_{G\otimes G} : G \otimes G \to G.$$

Proposition 2.4 *The map $\overline{\nu}_Y([D(z; r_1, r_2)])\big|_{G\otimes G}$ is continuous.*

Proof. From the definition of $\overline{\nu}_Y([D(z; r_1, r_2)])\big|_{G\otimes G}$, we see that for any $k, l \ge 0$,

$$\overline{\nu}_Y([D(z; r_1, r_2)]))(G_k \otimes G_l)$$

is in G_{k+l+1}. To prove that $\overline{\nu}_Y([D(z; r_1, r_2)])\big|_{G\otimes G}$ is continuous, we need only prove that for any $k, l \ge 0$, it is continuous as a map from $G_k \otimes G_l$ to G_{k+l+1}. Since the topology on G_k and G_l are defined to be the quotient topologies on

$$(X^{\otimes(k+1)} \otimes F_k^*)/(e_k|_{X^{\otimes(k+1)}\otimes F_k^*})^{-1}(0)$$

and

$$(X^{\otimes(l+1)} \otimes F_l^*)/(e_l|_{X^{\otimes(l+1)} \otimes F_l^*})^{-1}(0),$$

respectively, we need only prove that the map

$$\begin{aligned}(\overline{\nu}_Y([D(z; r_1, r_2)])) \circ (e_k|_{X^{\otimes(k+1)} \otimes F_k^*} \otimes e_l|_{X^{\otimes(l+1)} \otimes F_l^*}) :& \\ (X^{\otimes(k+1)} \otimes F_k^*) \otimes (X^{\otimes(l+1)} \otimes F_l^*) &\to G_{k+l+1}\end{aligned}$$

is continuous. On the other hand, by definition,

$$\begin{aligned}&(\overline{\nu}_Y([D(z; r_1, r_2)])) \circ (e_k|_{X^{\otimes(k+1)} \otimes F_k^*} \otimes e_l|_{X^{\otimes(l+1)} \otimes F_l^*}) \\ &\quad = e_{k+l+1}|_{X^{\otimes(k+l+2)} \otimes F_{k+l+1}^*} \circ (I_{X^{\otimes(k+l+1)}} \otimes \beta_{k,l}) \circ \sigma_{23},\end{aligned}$$

where $I_{X^{\otimes(k+l+1)}}$ is the identity map on $X^{\otimes(k+l+1)}$ and

$$\sigma_{23} : (X^{\otimes(k+1)} \otimes F_k^*) \otimes (X^{\otimes(l+1)} \otimes F_l^*) \to X^{\otimes(k+l+2)} \otimes F_k^* \otimes F_l^*$$

is the map permuting the second and the third tensor factor, that is,

$$\sigma_{23}(\mathcal{X}_1 \otimes \mu \otimes \mathcal{X}_2 \otimes \nu) = \mathcal{X}_1 \otimes \mathcal{X}_2 \otimes \mu \otimes \nu$$

for $\mathcal{X}_1 \in X^{\otimes(k+1)}$, $\mu \in F_k^*$, $\mathcal{X}_2 \in X^{\otimes(l+1)}$ and $\nu \in F_l^*$. Since the topology on G_{k+l+1} is defined to be the quotient topology on

$$(X^{\otimes(k+l+2)} \otimes F_{k+l+1}^*)/(e_{k+l+1}|_{X^{\otimes(k+l+2)} \otimes F_{k+l+1}^*})^{-1}(0),$$

we need only prove that the map $(I_{X^{\otimes(k+l+1)}} \otimes \beta_{k,l}) \circ \sigma_{23}$ is continuous. But $I_{X^{\otimes(k+l+1)}}$ and σ_{23} are obviously continuous and $\beta_{k,l}$ is continuous by Proposition 2.3. So $(I_{X^{\otimes(k+l+1)}} \otimes \beta_{k,l}) \circ \sigma_{23}$ is indeed continuous. □

Since G is dense in H, we can extend $(\overline{\nu}_Y([D(z; r_1, r_2)]))\big|_{G \otimes G}$ to a linear map $\overline{\nu}_Y([D(z; r_1, r_2)])$ from $H \widetilde{\otimes} H$ to H.

Theorem 2.5 *The map $\overline{\nu}_Y([D(z; r_1, r_2)])$ is a continuous extension of*

$$\nu_Y((z; \mathbf{0}, (1/r_1, \mathbf{0}), (1/r_2, \mathbf{0})))$$

to $H \widetilde{\otimes} H$. That is, $\overline{\nu}_Y([D(z; r_1, r_2)])$ is continuous and

$$\overline{\nu}_Y([D(z; r_1, r_2)])\big|_{V \otimes V} = \nu_Y((z; \mathbf{0}, (1/r_1, \mathbf{0}), (1/r_2, \mathbf{0}))).$$

Proof. The continuity of $\overline{\nu}_Y([D(z; r_1, r_2)])$ follows from the definition and Proposition 2.4.

For any $m_1, \dots, m_k \in \mathbb{Z}$, let $\mu_{m_1,\dots,m_k}$ be an element of F_k^* defined by

$$\begin{aligned}
&\mu_{m_1,\dots,m_k}(g_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes u))\\
&\quad = \frac{1}{2\pi\sqrt{-1}} \oint_{|z_1|=\epsilon_1} \cdots \frac{1}{2\pi\sqrt{-1}} \oint_{|z_k|=\epsilon_k} z_1^{m_1} \cdots z_k^{m_k} \cdot\\
&\qquad\qquad \cdot g_k(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes u) dz_k \cdots dz_k
\end{aligned}$$

for $\lambda \in \tilde{G}$, $u_1, \dots, u_k, u \in V$, where $\epsilon_1, \dots, \epsilon_k$ are arbitrary positive real numbers satisfying $\epsilon_1 > \cdots > \epsilon_k$. Since V is generated by X, elements of the form

$$e_k(u_1 \otimes \cdots \otimes u_k \otimes u \otimes \mu_{m_1,\dots,m_k}),$$

$k \geq 0$, $u_1, \dots, u_k, u \in X$ and $m_1, \dots, m_k \in \mathbb{Z}$, span V.

Let

$$h_1 = e_k(u_1 \otimes \cdots \otimes u_k \otimes u \otimes \mu_{m_1,\dots,m_k})$$

and

$$h_2 = e_l(v_1 \otimes \cdots \otimes v_l \otimes v \otimes \mu_{n_1,\dots,n_l})$$

be two such elements of V. Then

$$\begin{aligned}
&\langle \lambda, (\overline{\nu}_Y([D(z; r_1, r_2)]))(h_1 \otimes h_2)\rangle\\
&\quad = \langle \lambda, e_{k+l+1}(u_1 \otimes \cdots \otimes u_k \otimes u\\
&\qquad\qquad \otimes v_1 \otimes \cdots \otimes v_l \otimes v \otimes \beta_{k,l}(\mu_{m_1,\dots,m_k}, \mu_{n_1,\dots,n_l}))\rangle\\
&\quad = \beta_{k,l}(\mu_{m_1,\dots,m_k}, \mu_{n_1,\dots,n_l})\\
&\qquad\qquad (g_{k+l+1}(\lambda \otimes u_1 \otimes \cdots \otimes u_k \otimes u \otimes v_1 \otimes \cdots \otimes v_l \otimes v))\\
&\quad = \mu_{m_1,\dots,m_k}(g_k(\alpha_l(\lambda \otimes v_1 \otimes \cdots \otimes v_l \otimes v \otimes \mu_{n_1,\dots,n_l}) \otimes u_1 \otimes \cdots \otimes u_k \otimes u)).
\end{aligned} \tag{2.15}$$

As in the proof of Proposition 2.3, to avoid notational confusions, we view

$$g_k(\alpha_l(\lambda \otimes v_1 \otimes \cdots \otimes v_l \otimes v \otimes \mu_{n_1,\dots,n_l}) \otimes u_1 \otimes \cdots \otimes u_k \otimes u)$$

as an element of F_k^{ζ} and

$$g_l((P_n(Q(u_1, \dots, u_k, v; \zeta_1, \dots, \zeta_k)) \diamond_{[D(z;r_1,r_2)]} \lambda) \otimes v_1 \otimes \cdots \otimes v_l \otimes v)$$

as an element of F_l^η. We also view $\mu_{m_1,\dots,m_k}$ and $\nu_{n_1,\dots,n_l}$ as elements of $(F_k^\zeta)^*$ and $(F_l^\eta)^*$, respectively. Then the right-hand side of (2.15) is equal to

$$
\begin{aligned}
&\mu_{m_1,\dots,m_k}\Bigg(\sum_{n\in\mathbb{Z}}\alpha_l(\lambda\otimes v_1\otimes\cdots\otimes v_l\otimes v\otimes\mu_{n_1,\dots,n_l})\\
&\qquad\qquad (P_n(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_k))))\Bigg)\\
&\quad=\mu_{m_1,\dots,m_k}\Bigg(\sum_{n\in\mathbb{Z}}\mu_{n_1,\dots,n_l}(g_l((P_n(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_k))\diamond_{[D(z;r_1,r_2)]}\lambda)\\
&\qquad\qquad \otimes v_1\otimes\cdots\otimes v_l\otimes v))\Bigg)\\
&\quad=\mu_{m_1,\dots,m_k}\Bigg(\sum_{n\in\mathbb{Z}}\mu_{n_1,\dots,n_l}\Bigg(\sum_{m\in\mathbb{Z}}(P_n(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_k))\diamond_{[D(z;r_1,r_2)]}\lambda)\\
&\qquad\qquad (P_m(Q(v_1,\dots,v_l,v;\eta_1,\dots,\eta_l)))\Bigg)\Bigg)\\
&\quad=\mu_{m_1,\dots,m_k}\Bigg(\sum_{n\in\mathbb{Z}}\mu_{n_1,\dots,n_l}\Bigg(\sum_{m\in\mathbb{Z}}\sum_{p\in\mathbb{Z}}\lambda(P_p(r_1^{L(0)}\cdot\\
&\qquad\qquad \cdot Y(P_n(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_k)),z)\cdot\\
&\qquad\qquad \cdot r_2^{L(0)}P_m(Q(v_1,\dots,v_l,v;\eta_1,\dots,\eta_l)))\Bigg)\Bigg)\\
&\quad=\frac{1}{2\pi\sqrt{-1}}\oint_{|\zeta_1|=\epsilon_1}\cdots\frac{1}{2\pi\sqrt{-1}}\oint_{|\zeta_k|=\epsilon_k}\zeta_1^{m_1}\cdots\zeta_k^{m_k}\cdot\\
&\qquad\qquad \cdot\sum_{n\in\mathbb{Z}}\frac{1}{2\pi\sqrt{-1}}\oint_{|\eta_1|=\delta_1}\cdots\frac{1}{2\pi\sqrt{-1}}\oint_{|\eta_l|=\delta_l}\eta_1^{n_1}\cdots\eta_l^{n_l}\cdot\\
&\qquad\qquad \cdot\sum_{m\in\mathbb{Z}}\sum_{p\in\mathbb{Z}}\lambda(P_p(r_1^{L(0)}Y(P_n(Q(u_1,\dots,u_k,u;\zeta_1,\dots,\zeta_k)),z)\cdot\\
&\qquad\qquad \cdot r_2^{L(0)}P_m(Q(v_1,\dots,v_l,v;\eta_1,\dots,\eta_l)))\\
&\quad=\sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}}\sum_{p\in\mathbb{Z}}\lambda(P_p(r_1^{L(0)}Y(P_n(\mathrm{Res}_{x_1}\cdots\mathrm{Res}_{x_k}x_1^{m_1}\cdots x_k^{m_k}\cdot\\
&\qquad\qquad \cdot Y(u_1,x_1)\cdots Y(u_k,x_1)u),z)r_2^{L(0)}\cdot\\
&\qquad\qquad \cdot P_n(\mathrm{Res}_{y_1}\cdots\mathrm{Res}_{y_l}y_1^{n_1}\cdots y_l^{n_l}Y(v_1,y_1)\cdots Y(v_l,y_l)v))\\
&\quad=\sum_{p\in\mathbb{Z}}\lambda(P_p(Y(r_1^{L(0)}h_1,z)r_2^{L(0)}h_2))
\end{aligned}
$$

$$= \langle \lambda, Y(r_1^{L(0)}h_1, z)r_2^{L(0)}h_2\rangle. \tag{2.16}$$

From (2.15) and (2.16), we get

$$\begin{aligned}
&(\overline{\nu}_Y([D(z; r_1, r_2)]))(h_1 \otimes h_2)\\
&\quad = Y(r_1^{L(0)}h_1, z)r_2^{L(0)}h_2\\
&\quad = (\nu_Y((z; \mathbf{0}, (1/r_1, \mathbf{0}), (1/r_2, \mathbf{0}))))(h_1 \otimes h_2),
\end{aligned}$$

proving the theorem. □

3 A locally convex completion of a finitely-generated module and the vertex operator map

In this section, we discuss modules. Since the constructions and proofs are all similar to the case of algebras, we shall only state the results and point out the slight differences between the constructions and proofs in the case of modules and those in the case of algebras.

Let V be a $\mathbb{Z}$-graded finitely generated vertex algebra satisfying the standard grading-restriction axioms and W a $\mathbb{C}$-graded finitely-generated V-module satisfying the standard grading-restriction axioms, that is,

$$W = \coprod_{n\in\mathbb{C}} W_{(n)},$$

$$\dim W_{(n)} < \infty,$$

$n \in \mathbb{C}$ and

$$W_{(n)} = 0$$

for $n \in \mathbb{C}$ whose real part is sufficiently small. Let M be the finite-dimensional vector space spanned by a set of generators of W. As in Section 2, we assume that X is a finite-dimensional subspace containing $\mathbf{1}$ of V spanned by a set of generators of V.

We construct spaces $\tilde{G}^W$ and F_k^W, $k \geq 0$, and their duals in the same way as the constructions of $\tilde{G}$ and F_k, $k \geq 0$, and their duals in Section 2, except that V^* is replaced by W^*, $v \in V$ is replaced by $w \in W$ and the vertex operator map Y for V are replaced by vertex operator map Y_W for the module W. We also define linear maps g_k^W, $\iota_{F_k^W}$, γ_k^W, e_k^W, $k \geq 0$, and linear maps induced from them in the same way as in the definitions of g_k, ι_{F_k}, γ_k, e_k, $k \geq 0$, and induced linear maps in Section 2, except that the spaces are replaced by the corresponding spaces involving W and M.

For $k \geq 0$, e_k^W is a linear map from $V^{\otimes k} \otimes W \otimes (F_k^W)^*$ to $(\tilde{G}^W)^*$. Let

$$G_k^W = e_k^W(X^{\otimes k} \otimes M \otimes (F_k^W)^*)$$

and

$$G^W = \bigcup_{k \geq 0} G_k^W.$$

As in the case of $\tilde{G}$ and H, the spaces $\tilde{G}^W$ and $(\tilde{G}^W)^*$ form a dual pair of vector spaces and thus we have a locally convex topology on $(\tilde{G}^W)^*$. As in Section 2, we have:

Proposition 3.1 *For any $k \geq 0$, the linear map*

$$e_k^W|_{X^{\otimes k} \otimes M \otimes (F_k^W)^*} : X^{\otimes k} \otimes M \otimes (F_k^W)^* \to (\tilde{G}^W)^*$$

is continuous. In particular,

$$X^{\otimes k} \otimes M \otimes (F_k^W)^* / (e_k^W|_{X^{\otimes k} \otimes M \otimes (F_k^W)^*})^{-1}(0)$$

and G_k^W is a locally convex space. □

For $k \geq 0$, let H_k^W be the completion of G_k^W and

$$H^W = \bigcup_{k \geq 0} H_k^W.$$

Then H_k^W, $k \geq 0$, are complete locally convex space.

Proposition 3.2 *For $k \geq 0$, H_k^W can be embedded canonically in H_{k+1}^W and the topology on H_k^W is induced from the topology on H_{k+1}^W.* □

Theorem 3.3 *The vector space H^W equipped with the strict inductive limit topology is a locally convex completion of W. In particular, G^W is dense in H^W.* □

Next we extend the map

$$Y_W(r_1^{L(0)}\cdot, z)r_2^{L(0)}\cdot : V \otimes W \to \overline{W}$$

associated to $[D(z; r_1, r_2)]$ to a linear map $\overline{\nu}_{Y_W}([D(z; r_1, r_2)])$ from $H\tilde{\otimes}H^W$ to H^W.

We define $u \diamond_{[D(z;r_1,r_2)]} \lambda \in W^*$ for $\lambda \in \tilde{G}^W$ and $u \in V$, the maps

$$\alpha_l^W : \tilde{G}^W \otimes X^l \otimes W \otimes (F_l^W)^* \to \tilde{G},$$

$l \geq 0$, and

$$\beta_{k,l}^W : F_k^* \otimes (F_l^W)^* \to (F_{k+l+1}^W)^*,$$

$k, l \geq 0$, in the same way as in the definitions in Section 3 of $u \diamond_{[D(z;r_1,r_2)]} \lambda \in V^*$ for $\lambda \in \tilde{G}$ and $u \in V$, the maps

$$\alpha_l : \tilde{G}^W \otimes X^{l+1} \otimes F_l^* \to V^*,$$

$l \geq 0$, and

$$\beta_{k,l} : F_k^* \otimes F_l^* \to F_{k+l+1}^*,$$

$k, l \geq 0$. (Note that the image of α_l^W is in $\tilde{G}$, not in $\tilde{G}^W$. This is the reason why the domain of $\beta_{k,l}^W$ is $F_k^* \otimes (F_l^W)^*$, not $(F_k^W)^* \otimes (F_l^W)^*$.)

Let

$$h_1 = e_k(u_1 \otimes \cdots \otimes u_k \otimes u \otimes \mu) \in G_k$$

and

$$h_2 = e_l^W(v_1 \otimes \cdots \otimes v_l \otimes w \otimes \nu) \in G_l^W$$

where $u_1, \dots, u_k, u, v_1, \dots, v_l \in X$, $w \in M$ and $\mu \in F_k^*$, $\nu \in (F_l^W)^*$. Define

$$\begin{aligned}(\overline{\nu}_{Y_W}([D(z; r_0, r_1, r_2)]))(h_1 \otimes h_2) \\ = e_{k+l+1}^W(u_1 \otimes \cdots \otimes u_k \otimes u \otimes v_1 \otimes \cdots \otimes v_l \otimes w \otimes \beta_{k,l}^W(\mu, \nu)).\end{aligned}$$

Note that any element of G_k or G_l^W is a linear combination of elements of the form h_1 or h_2, respectively, above, and that k and l are arbitrary. Thus we obtain a linear map

$$\overline{\nu}_{Y_W}([D(z; r_0, r_1, r_2)])\big|_{G \otimes G^W} : G \otimes G^W \to G^W.$$

Proposition 3.4 *The map* $\overline{\nu}_{Y_W}([D(z; r_1, r_2)])\big|_{G \otimes G^W}$ *is continuous.* $\square$

By this proposition and the fact that G and G^W are dense in H and H^W, respectivel, we can extend $\overline{\nu}_{Y_W}([D(z;r_1,r_2)])\big|_{G\otimes G^W}$ to a continuous linear map $\overline{\nu}_{Y_W}([D(z;r_1,r_2)])$ from $H\tilde{\otimes}H^W$ to H^W.

Theorem 3.5 *The map $\overline{\nu}_{Y_W}([D(z;r_1,r_2)])$ is a continuous extension of*

$$Y_W(r_1^{L(0)}\cdot, z)r_2^{L(0)}\cdot : V\otimes W\to \overline{W}$$

to $H\tilde{\otimes}H^W$. $\square$

References

[B] R. E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Natl. Acad. Sci. USA* **83** (1986), 3068–3071.

[F] I. B. Frenkel, talk presented at the Institute for Advanced Study, 1988; and private communications.

[FHL] I. B. Frenkel, Y.-Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, preprint, 1989; *Memoirs Amer. Math. Soc.* **104**, 1993.

[FLM] I. B. Frenkel, J. Lepowsky, and A. Meurman, *Vertex operator algebras and the Monster*, Pure and Appl. Math., **134**, Academic Press, New York, 1988.

[H1] Y.-Z. Huang, *On the geometric interpretation of vertex operator algebras*, Ph.D thesis, Rutgers University, 1990.

[H2] Y.-Z. Huang, Geometric interpretation of vertex operator algebras, *Proc. Natl. Acad. Sci. USA* **88** (1991), 9964–9968.

[H3] Y.-Z. Huang, Applications of the geometric interpretation of vertex operator algebras, *Proc. 20th International Conference on Differential Geometric Methods in Theoretical Physics, New York, 1991*, ed. S. Catto and A. Rocha, World Scientific, Singapore, 1992, 333–343.

[H4] Y.-Z. Huang, Vertex operator algebras and conformal field theory, *Intl. Jour. of Mod. Phys* **A7** (1992), 2109–2151.

[H5] Y.-Z. Huang, Intertwining operator algebras, genus-zero modular functors and genus-zero conformal field theories, in: *Operads: Proceedings of Renaissance Conferences*, ed. J.-L. Loday, J. Stasheff, and A. A. Voronov, Contemporary Math., **202**, Amer. Math. Soc., Providence, 1997, 335–355.

[H6] Y.-Z. Huang, *Two-dimensional conformal geometry and vertex operator algebras*, Progress in Mathematics, Vol. 148, Birkhäuser, Boston, 1997.

[H7] Y.-Z. Huang, Genus-zero modular functors and intertwining operator algebras, *Internat. J. Math.* **9** (1998).

[K1] G. Köthe, *Topological vector spaces I*, Grundlehren der mathematischen Wissenschaften 159, English Edition, Springer-Verlag, New York, 1969.

[K2] G. Köthe, *Topological vector spaces II*, Grundlehren der mathematischen Wissenschaften 237, Springer-Verlag, New York, 1979.

[S1] G. B. Segal, The definition of conformal field theory, preprint, 1988.

[S2] G. B. Segal, Two-dimensional conformal field theories and modular functors, in: *Proceedings of the IXth International Congress on Mathematical Physics, Swansea, 1988*, Hilger, Bristol, 1989, 22–37.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, 110 FRELINGHUYSEN RD., PISCATAWAY, NJ 08854-8019

E-mail address: yzhuang@math.rutgers.edu